

Intravenous Pyelogram (IVP)

Patient Information Sheet No. SCR3

Your appointment:

Date:

Time:

Location:

.....
Please see our website or the back of this form for more branch details

Duration of Examination:

The examination will take between 30-60 minutes. Some kidneys empty at slower rates which may make the examination last up to 4 hours.

Please bring with you:

- Your request form
- All previous relevant imaging
- Medicare and any Healthcare cards

Payment on the day of the examination is requested.

For more information on this procedure please call one of our branches.

For more information regarding Benson Radiology please visit:
www.bensonradiology.com.au

What is an Intravenous Pyelogram?

An intravenous pyelogram (IVP) is an x-ray examination of the urinary tract (kidneys, ureters and urinary bladder). Iodine-based contrast media is injected intravenously. The contrast collects in the urinary tract, defining these areas by making them opaque on x-rays. This allows the radiologist (specialist medical imaging doctor) to visualise the anatomy and function of the kidneys and urinary tract and assess abnormalities in the urinary system.

What preparation is required?

A strong laxative must be taken the night before your appointment to reduce bowel material which may obscure clear imaging of your kidneys. You must have nothing to eat for two hours prior to the examination but you may drink a little water.

What will happen during the examination?

You will be asked to change into a gown and lie on the examination table. IVP equipment is specialised and is not available at all of our rooms. Preliminary x-rays of your urinary tract are then taken.

An injection of contrast will be given into a vein in your arm and the radiographer will begin taking x-rays of your kidneys. Please try to remain very still during the examination. You will be asked to hold your breath for a few seconds reducing the possibility of blurred images.

As the contrast material is processed by the kidneys a series of images is taken to determine the actual size of the kidneys and to capture the urinary tract in action as it begins to empty. We may apply a compression band around your body to better visualize the urinary structures leading from the kidney.

During the imaging process, you may be asked to turn from side to side and to hold several different positions to enable the radiologist to capture views from several angles. Near the end of the examination, you may be asked to empty your bladder so that an additional x-ray can be taken of your empty urinary bladder.

Are there any risks?

You will receive a small dose of x-ray radiation. The benefits of detecting disease are believed to outweigh any potential risks from receiving such a small dose. **Please, however, advise the radiographer if you are, or think you may be, pregnant.**

Problems related to the contrast injection are very rare. The injection will be discussed prior to the examination and you will be asked to complete a questionnaire and to give your consent. Most people experience a temporary warming sensation when the contrast is administered. This will dissipate quickly. **Please inform us if you are taking any diabetic medication, have kidney disease or have had a previous reaction to x-ray contrast.**

After the examination

At the completion of the IVP the needle will be removed. You should be able to resume your normal activities.

The images will be reviewed and reported by our radiologist. This can take up to 90 minutes. For your convenience we can generally deliver the imaging and report to your doctor by the next working day. Alternatively your doctor may request that you wait or return later to collect the imaging and report.

It is important you understand why and how we will be assisting you. If you have any questions, please call one of our branches or speak to our staff on the day of your visit.